

# WENN'S NICHT WEITERGEHT – STILLSTAND ODER PAUSE?

Hebammenforum Geburt KOMPAKT 2/2022: 12–14

- 1 Bundesamt für Gesundheit (BAG): Kaiserschnittgeburten in der Schweiz. 2015. <https://www.bag.admin.ch/bag/de/home/suche.html#kaiserschnitt> (Zugriff 29.1.21)
- 2 gynécologie suisse SGGG: Guideline Sectio Caesarea. 2015. <https://www.sggg.ch/fachthemen/leitlinien-sggg-dggg-oeggg/> (Zugriff 29.1.21)
- 3 Caughey AB, Cahill AG, Guise JM et al.: Safe prevention of the primary cesarean delivery. Am J Obstet Gynecol 2014; 210: 179–193.
- 4 Miller S, Abalos E, Chamillard M et al.: Beyond too little, too late and too much, too soon: a pathway towards evidence-based, respectful maternity care worldwide. In: The Lancet 2016; 388: 2176–2192.
- 5 World Health Organisation: WHO recommendations: Intrapartum care for positive childbirth experience. Genf 2018. <https://www.who.int/reproductive-health/publications/intrapartum-care-guidelines/en/> (Zugriff 29.1.21).
- 6 Hautakangas T, Palomaki O, Eidsto K et al.: Impact of obesity and other risk factors on labor dystocia in term primiparous women: A case control study. In: BMC Pregnancy Childbirth 2018; 18(1).
- 7 Kjærgaard H, Dykes A.K, Ottesen B et al.: Risk indicators for dystocia in low-risk nulliparous women: A study on lifestyle and anthropometrical factors. In: Journal of Obstetrics & Gynaecology Research 2010; 30(1), 25–29.
- 8 Selin L, Wallin G, Berg M: Dystocia in labour: Risk factors, management and outcome: A retrospective observational study in a Swedish setting. In: Acta Obstetricia et Gynecologica Scandinavica 2007; 87(2), 216–221.
- 9 Kjærgaard H, Olsen J, Ottesen B et al.: Obstetric risk indicators for labour dystocia in nulliparous women: A multi-centre cohort study. In: BMC Pregnancy Childbirth 2008; 8, 45.
- 10 Ragusa A, Gizzo S, Noventa M et al.: Prevention of primary caesarean delivery: Comprehensive management of dystocia in nulliparous patients at term. In: Archives of Gynecology and Obstetrics 2015; 294(4), 753–761.
- 11 Oladapo OT, Diaz V, Bonet M et al.: Cervical dilatation patterns of »low-risk« women with spontaneous labour and normal perinatal outcomes: A systematic review. In: BJOG 2018; 125: 944–954.
- 12 Fischer-Dückelmann A: Die Frau als Hausärztin. Ein ärztliches Nachschlagebuch für die Frau. Stuttgart 1905: Süddeutsches Verlags-Institut.
- 13 American College of Obstetricians and Gynecologists: Committee Opinion No. 650: Physical Activity and Exercise During Pregnancy and the Postpartum Period. In: Obstetrics & Gynecology 2015; e135.
- 14 Carlson NS, Hernandez TL, Hurt JK: Parturition dysfunction in obesity: time to target the pathobiology. In: Reproductive Biology and Endocrinology 2015; 13:135.
- 15 Konje JC, Oladapi AL: Nutrition and obstructed labor. In: American Journal of Clinical Nutrition 2000; 72(1), 291–297.
- 16 Neal JL, Lamp JM, Buck JS: Outcomes of nulliparous women with spontaneous labour onset admitted to hospitals in preactive versus active labour. In: Journal of Midwifery & Women's Health 2014; 29: 28–34.
- 17 Bumm E: Grundriss zum Studium der Geburtshilfe in achtundzwanzig Vorlesungen und sechshundert-sechsundzwanzig bildlichen Darstellungen im Text und auf drei Tafeln. Springer-Verlag Berlin Heidelberg. 14. und 15. verbesserte Auflage 1922.
- 18 Stadelmann I: Die Hebammensprechstunde. Stadelmann Verlag. 8. korrig. Aufl. 2005.
- 19 Schmid V: Der Geburtsschmerz als endokriner Reizfaktor. In: Hebamme.ch 2007; 6.
- 20 Liao J, Buhimschi CS, Norwitz ER: Normal Labor: Mechanism and Duration. In: Obstetrics and Gynecology clinics of North America 2005; 32, 145–164.
- 21 Wei S, Wo BL, Qi HP et al: Early amniotomy and early oxytocin for prevention of, or therapy for, delay in first stage spontaneous labour compared with routine care. In: Cochrane Database of Systematic Reviews 2013; 8, Cd006794.
- 22 Arnon Z, Dor A, Bazak H et al.: Complementary medicine for labouring women: a qualitative study of the effects of reflexology. In: Journal of Complementary and Integrative Medicine 2018 Jul 19;16(1).
- 23 Sakala C: (1988) Content of care by independent midwives: assistance with pain in labor and birth. In: Social Science and Medicine 1988; 26(11):1141–1158.
- 24 Leap N, Sandall J, Buckland S et al.: Journey to Confidence: Women's Experiences of Pain in Labour and Relational Continuity of Care. In: Journal of Midwifery & Women's Health 2010; 55(3).
- 25 Sandall J, Soltani H, Gates S et al.: Midwife-led continuity models versus other models of care for childbearing women. In: Cochrane Database of Systematic Reviews 2016; Band 4, Artikelnr. CD004667.
- 26 Hodnett ED, Gates S, Hofmeyr G J et al.: Continuous support for women during childbirth. In: Cochrane Database of Systematic Reviews 2013; 7, 1–118.
- 27 National Institute for Health and Care Excellence: Intrapartum care for healthy women and babies. 2014. <https://www.nice.org.uk/guidance/cg190> (Zugriff 29.1.21).