

# SICHERHEIT DURCH FETALE ÜBERWACHUNG?

Hebammenforum 6/2022; 23: 8–14

- 1 Wilson J, Jungner G: Principles and practice of screening for disease. WHO Genf, 1968
- 2 World Health Organization (WHO): Vorsorgeuntersuchung und Screening: ein kurzer Leitfaden. Genf, 2020
- 3 Sartwelle TP, Johnston JC: Cerebral palsy litigation: change course or abandon ship. *J Child Neurol* 2015; 30(7): 828–41
- 4 Rei M, Ayres-de-Campos D, Bernardes J: Neurological damage arising from intrapartum hypoxia/acidosis. *Best Pract Res Clin Obstet Gynaecol* 2016; 30: 79–86.
- 5 Nelson KB, Leviton A: How much of neonatal encephalopathy is due to birth asphyxia? *Am J Dis Child* 1991; 145: 1325–31
- 6 Nelson KB, Sartwelle TP, Rouse DJ: Electronic fetal monitoring, cerebral palsy, and caesarean section: assumptions versus evidence. *BMJ (Online)* 2016. 1;355: i6405
- 7 Sartwelle TP: Electronic fetal monitoring: a bridge too far. *J Leg Med* 2012; 33(3):313–79
- 8 Lorain P, Bower A, Gottardi E et al.: Risk factors for hypoxic-ischemic encephalopathy in cases of severe acidosis: a case-control study. *Acta Obstet Gynecol Scand* 2022; 101(4): 471–78
- 9 Graham EM, Adami RR, McKenney SL et al.: Diagnostic accuracy of fetal heart rate monitoring in the identification of neonatal encephalopathy. *Obstet Gynecol* 2014; 124(3): 507–513
- 10 Farquhar CM, Armstrong S, Masson V, et al.: Clinician identification of birth asphyxia using intrapartum cardiotocography among neonates with and without encephalopathy in new zealand. *JAMA Netw Open* 2020. 3(2): e1921363
- 11 siehe 11
- 12 National Institute for Health and Care Excellence (NICE): Intrapartum care for healthy women and their babies. 2014; [www.nice.org.uk/guidance/cg190/resources/intrapartum-care-for-healthy-women-and-babies-pdf-35109866447557](http://www.nice.org.uk/guidance/cg190/resources/intrapartum-care-for-healthy-women-and-babies-pdf-35109866447557) (Zugriff 22.4.22)
- 13 Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften (AWMF): S3-Leitlinie 015-083 »Die vaginale Geburt am Termin«. 2021; [www.awmf.org/leitlinien/detail/ll/015-083.html](http://www.awmf.org/leitlinien/detail/ll/015-083.html) (Zugriff 22.4.22)
- 14 Sholapurkar SL: Intermittent auscultation (surveillance) of fetal heart rate in labor: a progressive evidence-backed approach with aim to improve methodology, reliability and safety. *J Matern Fetal Neonatal Med* 2020; 30: 1–7
- 15 Blix E, Maude R, Hals E et al.: Intermittent auscultation fetal monitoring during labour: a systematic scoping review to identify methods, effects, and accuracy. *PLoS One* 2019; 14(7): e0219573
- 16 Lewis D, Downe S: FIGO consensus guidelines on intrapartum fetal monitoring: Intermittent auscultation. *Int J Gynaecol Obstet* 2015; 131(1): 9–12
- 17 Smith V, Begley CM, Clarke M, Devane D: Professionals' views of fetal monitoring during labour: a systematic review and thematic analysis. *BMC Pregnancy Childbirth* 2012; 12: 166